

References

- Bates, D., M. Mächler, B. Bolker, and S. Walker (2014). Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823*. <http://arxiv.org/abs/1406.5823>.
- Belenky, G., N. Wesensten, D. Thorne, M. Thomas, H. Sing, D. Redmond, M. Russo, and T. Balkin (2003). Patterns of performance degradation and restoration during sleep restriction and subsequent recovery: a sleep dose-response study. *Journal of Sleep Research* 12(1), 1–12.
- Breslow, N. E. and D. G. Clayton (1993). Approximate Inference in Generalized Linear Mixed Models. *Journal of the American Statistical Association* 88(421), 9–25.
- Cleveland, W. S. (1979). Robust Locally Weighted Regression and Smoothing Scatterplots. *Journal of the American Statistical Association* 74(368), 829–836.
- Crainiceanu, C. and D. Ruppert (2004). Likelihood ratio tests in linear mixed models with one variance component. *Journal of the Royal Statistical Society. Series B* 66(1), 165–185.
- Dempster, A., N. Laird, and D. Rubin (1977). Maximum Likelihood from Incomplete Data via the EM Algorithm. *Journal of the Royal Statistical Society. Series B* 39(1), 1–38.
- Diggle, P. (1989). Testing for Random Dropouts in Repeated Measurement Data. *Biometrics* 45(4), 1255–1258.
- Diggle, P., P. Heagerty, K.-Y. Liang, and S. Zeger (2002). *Analysis of Longitudinal Data*. Oxford University Press.
- Fitzmaurice, G., M. Davidian, G. Verbeke, and G. Molenberghs (2008). *Longitudinal Data Analysis*. Chapman & Hall/CRC.
- Fitzmaurice, G., N. Laird, and J. Ware (2004). *Applied longitudinal analysis*. Wiley-Interscience.
- Greven, S. and T. Kneib (2010). On the Behaviour of Marginal and Conditional AIC in Linear Mixed Models. *Biometrika* 97(4), 773–789.

- Henderson, C. (1950). Estimation of genetic parameters. *Annals of Mathematical Statistics* 21, 309–310.
- Joe, H. (2008). Accuracy of Laplace approximation for discrete response mixed models. *Computational Statistics & Data Analysis* 52(12), 5066–5074.
- Lesaffre, E., M. Asefa, and G. Verbeke (1999). Assessing the goodness-of-fit of the Laird and Ware model – an example: the Jimma Infant Survival Differential Longitudinal Study. *Statistics in Medicine* 18, 835–854.
- Liang, K.-Y. and S. Zeger (1986). Longitudinal Data Analysis Using Generalized Linear Models. *Biometrika* 73(1), 13–22.
- Little, R. (2008). Selection and pattern-mixture models. In *Longitudinal Data Analysis*, pp. 409–432. Chapman & Hall/CRC.
- Little, R. and D. Rubin (1987). *Statistical Analysis with Missing Data*. J. Wiley & Sons.
- Molenberghs, G. and G. Verbeke (2005). *Models for Discrete Longitudinal Data*. Springer.
- Molenberghs, G. and G. Verbeke (2007). Likelihood ratio, score, and Wald tests in a constrained parameter space. *The American Statistician* 61(1), 22–27.
- Neuhaus, J., J. Kalbfleisch, and W. Hauck (1991). A Comparison of Cluster-Specific and Population-Averaged Approaches for Analyzing Correlated Binary Data. *International Statistical Review* 59(1), 25–35.
- Pinheiro, J. and D. Bates (2000). *Mixed-effects models in S and S-PLUS*. Springer.
- Potthoff, R. and S. Roy (1964). A Generalized Multivariate Analysis of Variance Model Useful Especially for Growth Curve Problems. *Biometrika* 51(3/4), 313–326.
- Robins, J., A. Rotnitzky, and L. Zhao (1995). Analysis of Semiparametric Regression Models for Repeated Outcomes in the Presence of Missing Data. *Journal of the American Statistical Association* 90(429), 106–121.

- Robinson, G. (1991). That BLUP is a good thing: the estimation of random effects. *Statistical Science* 6(1), 15–32.
- Royston, P. and D. G. Altman (1994). Regression Using Fractional Polynomials of Continuous Covariates: Parsimonious Parametric Modelling. *Journal of the Royal Statistical Society. Series C (Applied Statistics)* 43(3), 429–467.
- Rubin, D. (1976). Inference and Missing Data. *Biometrika* 63(3), 581–592.
- Rubin, D. (1987). *Multiple Imputation for Nonresponse in Surveys*. J. Wiley & Sons.
- Ruppert, D., P. Wand, and R. Carroll (2003). *Semiparametric Regression*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press.
- Self, S. and K. Liang (1987). Asymptotic properties of maximum likelihood estimators and likelihood ratio tests under nonstandard conditions. *Journal of the American Statistical Association* 82(398), 605–610.
- Silverman, B. W. (1985). Some aspects of the spline smoothing approach to non-parametric regression curve fitting. *Journal of the Royal Statistical Society. Series B*, 1–52.
- Stram, D. and J. Lee (1994). Variance Components Testing in the Longitudinal Mixed Effects Model. *Biometrics* 50(4), 1171–1177.
- Stram, D. and J. Lee (1995). Correction to 'Variance components testing in the longitudinal mixed effects model'. *Biometrics* 51(3), 1196.
- Swihart, B., B. Caffo, B. James, M. Strand, B. Schwartz, and N. Punjabi (2010). Lasagna plots: A saucy alternative to spaghetti plots. *Epidemiology* 21(5), 621–625.
- Townsend, Z., J. Buckley, and M. Harada (2013). The Choice between Fixed and Random Effects. In *The SAGE Handbook of Multilevel Modeling*, pp. 73–89. SAGE Publications Ltd.
- Vaida, F. and S. Blanchard (2005). Conditional Akaike information for mixed-effects models. *Biometrika* 92(2), 351–370.

- Verbeke, G. and G. Molenberghs (2000). *Linear Mixed Models for Longitudinal Data*. Springer.
- Wood, S. (2006). *Generalized additive models: an introduction with R*. Chapman & Hall/CRC.
- Zeger, S., K.-Y. Liang, and P. Albert (1988). Models for Longitudinal Data: A Generalized Estimating Equation Approach. *Biometrics* 44(4), 1049–1060.